

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/69243 A1

(51) International Patent Classification⁷: **G01N 33/497,**
A01C 1/02, C12M 1/34

[NL/NL]; Rhynggeesterstraatweg 52, NL-2341 BV Oegst-
geest (NL). **KÖNIG, Johan, Willem** [NL/NL]; Rosser 5,
NL-2201 VE Noordwijk (NL).

(21) International Application Number: **PCT/NL01/00217**

(22) International Filing Date: **16 March 2001 (16.03.2001)**

(74) Agent: **PRINS, A., W.**; Verenigde, Nieuwe Parklaan 97,
NL-2587 BN The Hague (NL).

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
00200990.0 **17 March 2000 (17.03.2000)** **EP**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **NED-
ERLANDSE ORGANISATIE VOOR TOEGEPAST-
NATUURWETENSCHAPPELIJK ONDERZOEK
TNO** [—/NL]; Schoemakerstraat 97, NL-2628 VK Delft
(NL).


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **VAN DUIJN, Albert**

[Continued on next page]

(54) Title: **MEASURING METABOLIC RATE CHANGES**



(57) Abstract: The invention relates to methods for measuring metabolic states or rates or changes therein, such as growth rates, dying rates, cell division, metabolite production, and other biological activities of organisms, in particular of small multi-cellular organisms, of seeds and seedlings and of micro-organisms, such as fungi, yeast, bacteria, plant or animal cells and cultures thereof. The invention provides a method for determining a change in metabolic rate of at least one organism comprising placing said organism or part thereof in a confined container and repeatedly or continually measuring the concentration of a metabolic gas in said confined container to determine changes in consumption or production of said gas by said organism wherein said gas concentration is determined without essentially affecting the concentration of said gas in said confined container.

WO 01/69243 A1



Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Title: Measuring metabolic rate changes.

The invention relates to methods for measuring metabolic states, metabolic rates and changes therein, such as growth rates, dying rates, cell division, metabolite production, and other biological activities of organisms, in particular of small multi-cellular organisms, of seeds and seedlings and of
5 micro-organisms, such as fungi, yeast, bacteria, plant or animal cells and cultures thereof.

Biological activities of organisms are manifold and are often studied by the chemical, physical, physiological or morphological ways they manifest themselves. Culturing organisms, be it micro-organisms in cell culture, plants,
10 or others, requires managing these biological activities, and managing these activities often requires measuring the underlying metabolic activities. As an example herein germination of seeds is discussed, however, the invention extends to culturing other organisms where similar approaches apply.

A new plant formed by sexual reproduction starts as an embryo within
15 the developing seed, which arises from the ovule. When mature, the seed is the means by which the new individual is dispersed, though frequently the ovary wall or even extrafloral organs remain in close association to form a more complex dispersal unit as in grasses and cereals. The seed, therefore, occupies a critical position in the life history of the higher plant. The success with
20 which the new individual is established -the time, the place, and the vigour of the young seedling- is largely determined by the physiological and biochemical features of the seed. Of key importance to this success are the seed's responses to the environment and the food reserves it contains, which are available to sustain the young plant in the early stages of growth before it becomes an
25 independent, autotrophic organism, able to use light energy. People also depend on these activities for almost all of their utilisation of plants.

Cultivation of most crop species depends on seed germination, though, of course, there are exceptions when propagation is carried out vegetatively. Moreover, seed such as those of cereals and legumes are themselves major food

sources whose importance lies in the storage reserves of protein, starch, and oil laid down during development and maturation.

In the scientific literature the term germination is often used loosely and sometimes incorrectly and so it is important to clarify its meaning.

5 Germination begins with water uptake by the seed (imbibition) and ends with the start of elongation by the embryonic axis, usually the radicle. It includes numerous events, e.g., protein hydration, subcellular structural changes, respiration, macromolecular syntheses, and cell elongation, none of which is itself unique to germination. But their combined effect is to transform an
10 organism having a dehydrated, resting metabolism into an organism having an active metabolism, culminating in growth.

Germination *sensu stricto* therefore does not include seedling growth, which commences when germination finishes. Hence, it is incorrect, for example, to equate germination with seedling emergence from soil since
15 germination will have ended sometime before the seedling is visible. Seed testers often refer to germination in this sense because their interests lie in monitoring the establishment of a vigorous plant of agronomic value. However, physiologists do not encourage such a definition of the term germination but in general acknowledge its widespread use by seed technologists. It would
20 however be preferable to find a more defined definition. Processes occurring in the nascent seedling, such as mobilisation of the major storage reserves, are also not part of germination: they are postgermination events.

A seed in which none of the germination processes is taking place is said to be quiescent. Quiescent seeds are resting organs, generally having a low
25 moisture content (5-15%) with metabolic activity almost at a standstill. A remarkable property of seeds is that they are able to survive in this state, often for many years, and subsequently resume a normal, high level of metabolism. For germination to occur quiescent seeds generally need only to be hydrated under conditions that encourage metabolism, e.g., a suitable
30 temperature and presence of oxygen.

Components of the germination process, however, may occur in a seed that does not achieve radicle emergence. Even when conditions are apparently favourable for germination so that imbibition, respiration, synthesis of nucleic acids and proteins, and a host of other metabolic events all proceed, culmination in cell elongation does not occur, for reasons that are still poorly understood; such a seed expresses dormancy. Seeds that are dispersed from the parent plant already containing a block to the completion of germination show primary dormancy. Sometimes, a block(s) to germination develops in hydrated, mature seeds when they experience certain environmental conditions, and such seeds show induced or secondary dormancy. Dormant seeds are converted into germinable seeds (i.e., dormancy is broken) by certain "priming" treatments such as a light stimulus or a period at low or alternating temperature which nullify the block to germination but which themselves are not needed for the duration of germination process.

The extent to which germination has progressed can be determined roughly, say by measuring water uptake or respiration, but these measurements give us only a very broad indication of what stage of the germination process has been reached. No universally useful biochemical marker of the progress of germination has been found. The only stage of germination that we can time fairly precisely is its termination. Emergence of the axis (usually the radicle) from the seed normally enables us to recognise when germination has gone to completion, though in those cases where the axis may grow before it penetrates through the surrounding tissues, the completion of germination can be determined as the time when a sustained rise in fresh weight begins.

We are generally interested in following the germination behaviour of large numbers of seeds, e.g., all the seeds produced by one plant or inflorescence, or all those collected in a soil sample, or all those subjected to certain experimental treatment. The degree to which germination has been completed in a population is usually expressed as a percentage, normally

determined at time intervals over the course of the germination period which can be expressed in so-called germination curves, about which some general points should be made. Germination curves are usually sigmoidal, a minority of the seeds in the population germinates early, then the germination
5 percentage increases more or less rapidly, and finally few late germinators emerge. The curves are often positively skewed because a greater percentage germinates in the first half of the germination period than in the second. But although the curves have the same general shape, important differences in behaviour between populations are evident. For example, curves often flatten
10 off when only a low percentage of the seeds has germinated, showing that this population has low germination capacity, i.e., the proportion of seeds capable of completing germination is low. Assuming that these seeds are viable, the behaviour of the population could be related to dormancy or to environmental conditions, such as temperature or light, which do not favour germination of
15 most of the seeds.

The shape of the curves also depends on the uniformity of the population, i.e., the degree of simultaneity or synchrony of germination. When a limited percentage of seeds succeeds in germinating fairly early, but the remainder begin to do so only after a delay the population seems to consist of
20 two discrete groups: the quick and the slow germinators. This example also illustrates the point that populations with the same germination capacity can differ in other respects.

Three respiratory pathways are assumed to be active in the imbibed seed: glycolysis, the pentose phosphate pathway, and the citric acid (Krebs or
25 tricarboxylic acid) cycle. Glycolysis, catalysed by cytoplasmic enzymes, operates under aerobic and anaerobic condition to produce pyruvate, but in the absence of O_2 this is reduced further to ethanol, plus CO_2 , or to lactic acid if no decarboxylation occurs. Anaerobic respiration, also called fermentation, produces only two ATP molecules per molecule of glucose respired, in contrast
30 to six ATPs produced during pyruvate formation under aerobic conditions. In

the presence of O₂, further utilisation of pyruvate occurs within the mitochondrion: oxidative decarboxylation of pyruvate produces acetyl-CoA, which is completely oxidised to CO₂ and water via the citric acid cycle to yield up to a further 30 ATP molecules per glucose molecule respired. The
5 generation of ATP molecules occurs during oxidative phosphorylation when electrons are transferred to molecular O₂ along an electron transport (redox) chain via a series of electron carriers (cytochromes) located on the inner membrane of the mitochondrion. An alternative pathway for electron transport, which does not involve cytochromes, may also operate in
10 mitochondria.

The pentose phosphate pathway is an important source of NADPH, which serves as a hydrogen and electron donor in reductive biosynthesis, especially of fatty acid. Intermediates in this pathway are starting compounds for various biosynthetic processes, e.g., synthesis of various aromatics and
15 perhaps nucleotides and nucleic acid. Moreover, complete oxidation of hexose via the pentose phosphate pathway and the citric acid cycle can yield up to 29 ATPs.

Respiration by mature "dry" seeds (usual moisture content: 10-15%) of course is extremely low when compared with developing or germinating seeds,
20 and often measurements are confounded by the presence of a contaminating microflora. When dry seeds are introduced to water, there is an immediate release of gas. This so-called "wetting burst" which may last for several minutes, is not related to respiration, but is the gas that is released from colloidal adsorption as water is imbibed. This gas is released also when dead
25 seeds or their contents, e.g., starch, are imbibed.

Keto acids (e.g.; α -ketoglutarate, pyruvate), which are important intermediates in respiratory pathways, are chemically unstable and may be absent from the dry seed. A very early metabolic event during imbibition, occurring within the first few minutes after water enters the cells, is their

reformation from amino acids by deamination and transamination reactions (e.g., of glutamic acid and alanine).

The consumption of O₂ by many seeds follows a basic pattern although the pattern of consumption by the embryo differs ultimately from that by storage tissues. Respiration is considered to involve three or four phases:

Phase 1. Initially there is a sharp increase in O₂ consumption, which can be attributed in part to the activation and hydration of mitochondrial enzymes involved in the citric acid cycle and electron transport chain. Respiration during this phase increases linearly with the extent of hydration of the tissue.

Phase 2. This is characterised by a lag in respiration as O₂ uptake is stabilised or increases only slowly. Hydration of the seed parts is now completed and all pre-existing enzymes are activated. Presumably there is little further increase in respiratory enzymes or in the number of mitochondria during this phase. The lag phase in some seeds may occur in part because the coats or other surrounding structures limit O₂ uptake to the imbibed embryo or storage tissues, leading temporarily to partially anaerobic conditions. Removal of the testa from imbibed pea seeds, for example, diminishes the lag phase appreciably. Another possible reason for this lag is that the activation of the glycolytic pathway during germination is more rapid than the development of mitochondria. This could lead to an accumulation of pyruvate because of deficiencies in the citric acid cycle or oxidative phosphorylation (electron transport chain); hence, some pyruvate would be diverted temporarily to the fermentation pathway, which is not O₂ requiring.

Between phase 2 and 3 in the embryo the radicle penetrates the surrounding structures: germination is completed.

Phase 3. There is now a second respiratory burst. In the embryo, this can be attributed to an increase in activity of newly synthesised mitochondria and respiratory enzymes in the proliferating cells of the growing axis. The number of mitochondria in storage tissues also increases, often in association with the mobilisation of reserves. Another contributory factor of the rise in

respiration in both seeds parts could be an increased O₂ supply through the now punctured testa (or other surrounding structures).

Phase 4. This occurs only in storage tissues and coincides with their senescence following depletion of the stored reserves.

5 The lengths of phases 1-4 vary from species to species owing to such factors as differences in rates of imbibition, seed-coat permeability to oxygen, and metabolic rates. Moreover, the lengths of the phases will vary considerably with the ambient conditions, especially the temperature. In a few seeds, e.g., *Avena fatua*, there is no obvious lag phase in oxygen uptake. The
10 reasons for its absence are not known, but it could be because efficient respiratory systems become established early following imbibition, including the development of newly active mitochondria, thus ensuring a continued increase in O₂ utilisation. Also, coat impermeability might not restrict O₂ uptake prior to the completion of germination.

15 During germination a readily available supply of substrate for respiration must be present. This may be provided to a limited extent by hydrolysis of the major reserves, e.g., triacylglycerols, which are present in almost all parts of the embryo, including the radicle and hypocotyl, although their greatest concentration is in storage tissues. It is important to note,
20 however, that extensive mobilisation of reserves is a postgerminative event. Most dry seeds contain sucrose, and many contain one or more of the raffinose-series oligosaccharides: raffinose (galactosyl sucrose), stachyose (digalactosyl sucrose), and verbascose (trigalactosyl sucrose), although the latter is usually present only as a minor component. The distribution and amounts of these
25 sugars within seeds are very variable, even between different varieties of the same species.

During germination, sucrose and the raffinose-series oligosaccharides are hydrolysed, and in several species the activity of α -galactosidase, which cleaves the galactose units from the sucrose, increases as raffinose and
30 stachyose decline. Although there is little direct evidence that the released

monosaccharides are utilised as respiratory substrates, there is strong circumstantial evidence. Free fructose and glucose may accumulate in seeds during the hydrolysis of sucrose and the oligosaccharides, but there is no build-up of galactose (e.g., in mustard, *Sinapis alba*). Hence, it is probably rapidly
5 utilised, perhaps through incorporation into cell walls or into galactolipids of the newly forming membranes in the cells of developing seedling.

Virtually all metabolic pathways in living organisms, and not only those related to germination, relate to the uptake or release of metabolic gasses, of which the two most important are oxygen and carbon dioxide; examples of
10 others are carbon mono-oxide, nitric oxide, nitric dioxide, dinitric oxide, ethylene and ethanol. Classical is the way it could be demonstrated that oxygen is central to life. A mouse, placed under a glass bulb together with a burning candle, died when the flame dwindled and died, showing that also the mouse could not do without the oxygen. Undoubtedly, the level of carbon
15 dioxide in the glass bulb was, as a consequence, high.

The above example illustrates an archaic way of measuring the underlying metabolic activity of an organism. More modern methods have been developed which comprise measuring oxygen or other metabolic gasses in gas or liquid media. Oxygen, or other gasses, in gas are often measured by
20 analysis with gas-chromatography. In liquid gas contents are often measured by flushing some liquid through an electro-chemical measurement device.

For both types of measurements the sample is in general consumed and cannot be reused for other measurements. This has a number of further disadvantages: A container with the organism under study has to be opened
25 for a gas determination, which may disrupt the activities to be measured or otherwise hinder accurate determination. Also, for each point in a time series different samples are necessary for which the container has to be opened again. Normally this makes the number of samples very large and does not allow for using small containers to begin with. Furthermore, sample to sample
30 variability makes it very difficult to get reliable figures and the costs for

handling and measuring a sample are in general very high. The present invention recognizes this problem and provides a method for determining metabolic state or rate or a change therein of at least one organism or part thereof comprising placing said organism or part thereof in a confined
5 container and measuring the concentration of a metabolic gas in said confined container to determine consumption or production of said gas by said organism or part thereof wherein said gas concentration is determined without essentially affecting the concentration of said gas in said confined container. Such a method according to the invention has multiple advantages, for
10 example that the equilibrium of the gases within the confined container is not disturbed or influenced because the said container does not have to be opened to take a sample, thereby providing a very accurate and reliable method to determine the concentration of a metabolic gas in said confined container and as a consequence the metabolic state or rate or a change therein caused by at
15 least one organism or part thereof is accurate and reliably determined.

The invention provides a method for determining the metabolic state of at least one organism or part thereof comprising placing said organism or part thereof in a confined container and measuring the concentration of a metabolic
20 gas in said confined container to determine consumption or production of said gas by said organism or part thereof wherein said gas concentration is determined without essentially affecting the concentration of said gas in said confined container. If no change in metabolic gasses are detected (in practice for a sufficiently long period), it may for example be assumed that the
25 organism is dead or in a hibernating state, in particular now where the invention provides that no gas is consumed by measuring, all changes in gas concentration must thus be attributed to the production and/or consumption of a metabolic gas, thus of life, or at least in a state of life-like activity.

An example of an organism as disclosed herein within the experimental
30 part is a seed or a worm. It is clear to a person skilled in the art that different

organisms or parts thereof are tested by a method according to the invention as long as the organism or part thereof fits within a confined container. Therefor a method according to the invention is performed in a confined container which may have different sizes and/or shapes depending on the

5 organism or part thereof which need to be studied. An example of a part of an organism are the roots of a plant. The experimental part describes a rose from which the roots were put in a confined container. Another example of a part of an organism is a cell or a cell culture. Methods to arrive at a proper cell or cell culture are well known by the person skilled in the art. Preferably a method

10 according to the invention is used to determine changes in gas concentration of a predetermined organism or part thereof. Changes can therefor be attributed to a known, predetermined organism or part thereof.

Examples of metabolic gases from which the changes in concentration can be determined are oxygen, carbon dioxide, carbon mono-oxide, nitric oxide,

15 nitric dioxide, dinitric oxide, ethylene and ethanol. All these gases can be measured with different organo-metal complexes.

A confined container (also called confined space; the terms may be used interchangeably herein) is herein defined as a container that is properly shut to (essentially) avoid gas exchange between the confined container and the

20 surrounding and furthermore a confined container is defined as a container that is essentially not opened during measurements but to which additional substances (oxygen, nutrients, growth hormones, etc.) can be added with for example a valve or injection system. Because the container is essentially not opened all changes in a metabolic gas concentration are attributed to the

25 metabolic state or rate or a change therein of the organism or part thereof which is located in the container. A confined container has different shapes and/or sizes dependent on the organism or part thereof studied. It is clear to a person skilled in the art that after the organism or part thereof has been put in the container, the container is properly shut to (essentially) avoid gas

30 exchange between the confined container and the surrounding so that all

changes in gas concentration must be attributed to the production and/or consumption of a metabolic gas, thus of life, or at least in a state of life-like activity.

In a preferred embodiment, the invention provides a method for
5 determining a change in metabolic state or rate of at least one organism or part thereof comprising placing said organism or part thereof in a confined container and repeatedly or continually measuring the concentration of a metabolic gas in said container to determine changes in consumption or production of said gas by said organism or part thereof wherein said gas
10 concentration is determined without essentially affecting the concentration of said gas in said confined container. In one example of the invention one or more seeds are brought in a small confined container, along with some water to induce the germination process. Seeds can of course be totally immersed in water, which typically allows for measurements to be made in the liquid but
15 usually measurement of the air or gas above the seeds will be sufficient. Due to the germination at a certain point in time the seed(s) will start to consume oxygen and produce carbon dioxide. The oxygen concentration will drop from the moment the germination starts and the carbon dioxide concentration will rise. The gas concentration is preferably measured optically. This can for
20 example be achieved by a measuring device which is at least partly set up within the confined container, but measurements can also be made through a clear portion of the wall of the confined container, which for example could be made of glass.

In one embodiment, the invention provides an optical method based on
25 fluorescence quenching of fluorescent compounds by oxygen (1,2,3,4), to determine the oxygen levels inside a container, preferably without opening it. A sample can be measured over and over again in the time, and is not destroyed. Moreover, because the sample is not destroyed the number of samples necessary to do a time study is considerably lower compared to
30 conventional methods. In a preferred embodiment, the invention provides a

method wherein said gas concentration is determined by determining the fluorescence quenching of a fluorescent dye, preferably a suitable organo-metal, present in said confined container. For measuring oxygen, an oxygen sensitive dye such as a ruthenium bipyridyl complex, or Tris-Ru²⁺ 4,7 biphenyl 1,10 phenantrolin; or another Ru(ruthenium)-complex, or another organo-metal complex, such as an Os-complex or a Pt-complex, is suitable, for measuring carbon dioxide, or other gasses such as CO, NO, NO₂, N₂O, ethylene or ethanol, suitable sensitive organo-metal dyes, such as tris[2-(2-pyrazinyl)thiazole] ruthenium II (5) are used.

For example, the optical oxygen sensing measurement technique used herein is based on the fluorescence quenching of a metal organic fluorescent dye. The dye which is very sensitive to oxygen, is for example excited by a short laser light-pulse of for example 1 microsecond. After the excitation has stopped the oxygen sensitive dye emits fluorescent light with a decay curve which depends on the oxygen concentration. The process behind this phenomenon is called dynamic quenching.

Preferably said dye is present in a gas permeable compound such as silica or a hydrophobic polymer such as a (optionally fluoridated) silicone polymer, in PDMS (polydimethylsiloxane), in PTMSP (polytrimethylsilylpropyl), or in a mixture thereof but of course it can be contained in other suitable compounds as well. In a preferred embodiment the invention provides a method wherein said dye is present in at least a part of an inner coating of said confined container, for example situated on the inside of an optically transparent part of the confined container when measurement is from the outside.

Measuring can for example be achieved by measuring the fluorescence lifetime. The excited molecules are deactivated by oxygen in a collision process. The quenching process does not consume the gas (here the oxygen) so liquid medium does not necessarily have to be stirred to obtain the measurements. The fluorescence lifetime gets shorter because the probability of the molecules

to be deactivated gets higher for molecules which stay longer in the excited state. The effect is proportional with the quencher concentration. The relation between fluorescence lifetime and gas (here oxygen) concentration is given by the Stern Volmer equation (1)

5

$$\frac{\tau_0}{\tau} = 1 + C_{sv} \cdot [O_2]$$

10

where τ_0 is the fluorescence lifetime at quencher (O_2) concentration zero, τ is the fluorescence lifetime at a specific quencher (O_2) concentration. C_{sv} is the Stern-Volmer constant and $[O_2]$ is the gas concentration.

Measuring can also be achieved by measuring the fluorescence intensity.

15 The fluorescent compound is excited by a continuously radiating light source such as a LED and the fluorescence intensity is measured. More gas (here oxygen) caused less fluorescence. The relation between the oxygen concentration and the intensity is given by the Stern Volmer equation (2)

20

$$\frac{I_0}{I} = 1 + C_{sv} \cdot [O_2]$$

25

where I_0 is the fluorescence intensity at quencher (O_2) concentration zero, I is the fluorescence intensity at a specific quencher (O_2) concentration. C_{sv} is the Stern-Volmer constant and $[O_2]$ is the gas concentration.

Using the fluorescence lifetime method has the advantage that the
30 measurement is independent of the source intensity, detector efficiency,

fluorescent probe concentration etc. A method based on this principle is robust and less prone to drift. Moreover, because the quenching process does not consume oxygen or other metabolic gasses, the method as provided by the invention is very useful to measure metabolic rate changes of organisms by measuring an increase or decrease in metabolic gas production or consumption by said organism or organisms.

A method as provided by the invention is based on a time gated measurement (Fig.1). In this measurement method the fluorescence is determined in two time windows (A and B) after a light pulse. Fluorescence lifetime is a function of the ratio between A and B and is proportional to the oxygen concentration. The person skilled in the art is aware of the huge array of possible experimental set-ups. Figure 2 shows an example of a simplified experimental set-up. In this simplified set-up the confined container (having possibly different sizes and/or shapes) contains an oxygen sensitive coating situated on the inside of an optically transparent part of the confined container. Another possibility is to provide the oxygen sensitive substance to the material from which the confined container is made. Yet another possibility is to place the oxygen sensitive substance via a holder at any desired position within the confined container. The oxygen sensitive substance can be placed at every desired position as long as it is possible to reach the position with for example a laser to provoke excitation and to determine the fluorescence signal with a detector. Detection of the fluorescence is made visible by for example a measuring device or with help of a computer and suitable computer programs. In the above described simplified set-up the confined container is not physically part of the measuring device, but it is clear that it is possible to set-up a measuring device which is partly set-up in a confined container. A confined container can have different shapes and/or sizes and can be made of different materials as long as it is possible to perform measurements through a clear portion of the wall of the confined container, which for example could be made of glass. As described it is also possible that part of measuring device is part of

the confined container in which case it is not necessary for the wall of the container to be clear. Figure 3 shows a more detailed instrumental set up. A light source (e.g. LED or laser) is pulsed, the light pulses are filtered and excite the fluorescent dye located in the environment where the metabolic gas
5 has to be determined. The resulting fluorescence response is detected in a detector, the information is digitised, if needed the measurement is corrected (for temperature for example) and the gas concentration is calculated and displayed.

In the detailed description an example of a method according to the
10 invention is shown wherein said organism comprises a seed, and wherein said change in metabolic rate denotes germination. Oxygen consumption measurements on seeds during germination and priming are important for the following reasons. With regard to quality of seed batches (both for use as plant propagation method in e.g. horticulture and in industrial applications in e.g.
15 barley malting) the following aspects that can be achieved by measuring oxygen consumption during germination are important: (i) speed of germination, (ii) homogeneity of germination of a seed batch, (iii) monitoring system that is automated and (iv) possibility to measure large numbers of individual seeds. As the number of samples to be tested in seed companies is
20 very large and they are currently evaluated by eye, an automated system saves a lot of work. In addition many seeds should be kept in the dark during the test, the oxygen measurements can be performed in the dark in an automated system which solves the problems with the evaluation of these types of seeds. The homogeneity of seed batches is a quality aspect of prime
25 importance. This requires tests on individual seeds, so the possibility to automate the oxygen measurements in e.g. a measurement device using 96 wells plates offers an elegant solutions for the otherwise very labour-intensive test. During priming (this is a carefully controlled imbibition of seeds to obtain a pre-germination) it is important that the duration and extend of the priming
30 procedure is not too long (this results in primed seeds that cannot be dried

again). Monitoring of the metabolism (oxygen consumption) during the priming procedure will be an indicator of the progress of the priming that can be used to control the priming process. However, a method according to the invention is as well applicable to register a second respiratory burst as is often
5 identified in phase 3 of germination. The invention is furthermore used for quality assurance. For example seed batches primed or germinated by different methods or under different circumstances are tested. varieties are tested on for example their germination. To be able to perform high throughput screening a method according to the invention is preferably
10 miniaturised and/or automated. An example of such a automated/miniaturised device is depicted in Figure 7. It is clear to a person skilled in the art that, a preferably automated, quality assurance and/or high throughput screening is also used on another organism or part thereof. For example to test the effect of different kinds of insecticides on a mosquito. Of course, the invention provides
15 as well a method wherein said organism or part thereof comprises one or more micro-organisms or part thereof such as a protoplast, plastid (e.g. chloroplast) or mitochondrion, comprises plant cell cultures, comprises plant tissue explants, whole plants or seedlings, parts or organs of plants such as flowers, leaves, stems, roots, sexual organs, tubers, bulbs, fruits, or comprises animal
20 cell cultures, animal tissue explants, parts or organs of animals, blood, comprises a bacterium or bacterial cultures, a yeast cell or yeast cultures, a fungus or fungal cultures, and so on.

The invention provides a method wherein said change in metabolic rate denotes cell activity of said organism or part thereof or micro-organism or
25 cultures thereof, or, alternatively, wherein said change in metabolic rate denotes cell death, and to detect circumstances wherein such cell-activities thrive, or not. A method according to the invention is for example useful to detect (the onset of) sporulation of bacterial cultures, or microbial fermentation.

Within the field of seed technology, the invention provides a method to determine or monitor a rate of seed germination or development, to for example determine proper priming of seed batches. Therewith, the invention also provides a seed batch monitored with a method according to the invention.

5 Said seed batches have accurately been primed. Similarly, the invention provides a method to determine or monitor a rate of culture development of a cell- or tissue-culture comprising use of a method according to the invention and a cell- or tissue culture monitored with a method according to the invention. Other methods provides for example entail a method to determine

10 or monitor processing of waste water comprising use of a method according to the invention, or other processes where micro-biological fermentation plays a role.

As disclosed within the experimental part the invention is also used to determine the oxygen consumption of other organisms such as micro

15 organisms, animals, such as e.g. an insect or a worm. This part of the invention is e.g. useful to determine the presence of wood worms in a piece of (antique) wooden furniture or to determine the presence of wood worms in for example the wooden foundation or wooden floors in a house.

A method according to the invention is also used to test the effect of for

20 example an insecticide on its target by placing one or more targets in a confined space and determining the metabolic state or rate or a change therein and compare with one or more target(s) not treated with the insecticide. Preferably the tested and control targets have been selected on for example

25 their oxygen consumption, thereby providing good control experiments. Different analogues or derivatives of an insecticide are for example tested for their effectiveness.

The invention is further explained in the detailed description without limiting the invention thereto.

Detailed description

1. Determination of the start of seed germination by oxygen consumption

5

Most of conventional agriculture is engaged in growing plants from seeds. Plant breeding programs are dependent on the germination of the seeds obtained. Therefore the slow or no germination of seeds has a major impact on food production and research. The research on dormancy or environmental factors influencing the germination requires a simple method to monitor the germination process.

Methods.

15 One or more seeds (Triumph 1989, barley) are brought in a confined container, along with some water to induce the germination process. The container is closed. Due to the germination at a certain point in time the seed(s) will start to consume oxygen. Because the container is closed, the oxygen concentration will drop from the moment the germination starts. This can be monitored with
20 a special oxygen sensitive coating on the inside of an optically transparent part of the container. An advantage of optical oxygen determination is the fact that the coating itself does not consume any oxygen. In this way the start of the germination can be monitored accurately. Up to now only the appearance of a root was an indication of the germination. In the experiment the first root
25 showed after 10 to 14 hours. From the oxygen measurements we see that the germination activity showed after 3.5 hours. The oxygen consumption is an early indicator of seed germination.

The point where the germination starts can be calculated from the measured
30 oxygen levels, as for example given in table 1. The linear extrapolation of the

oxygen levels measured after 4 hours in the containers with 1, 2 and 3 seeds show an intersection with the oxygen level of an empty container at 3.5 hours after the addition of the water. This is the point in time where the metabolism of the germination starts. This is shown in figure 4. Figure 5 shows a general
5 course of oxygen levels with seeds in a container and figure 6 an example of a calculation. With this method it is possible to examine the effect of all kind of environmental influences on the germination process. It also gives an opportunity to influence the germination process in an early stage. This method is a simple and powerful tool in germination research and in the
10 priming of seeds.

Table 1 Oxygen content in μg of a sample container (approx. 1 ml) with a different number of seeds.

Hours	Container with 1 seed	Container with 2 seeds	Container with 3 seeds
0.0	284	284	284
0.5	263	271	261
1.5	274	253	261
2.5	284	279	267
3.5	274	261	270
4.5	270	259	266
5.5	279	256	223
6.5	277	253	227
7.5	271	231	204
8.5	282	257	198
9.5	269	227	160
11.5	275	202	125
14.0	262	172	51
24.0	220	41	1
29.0	250	14	7
32.0	238	9	4
35.5	226	8	4
38.0	201	8	4
50.0	171	6	5
53.0	191	8	5
61.0	159	9	4
72.0	135	7	3

2. **Oxygen consumption of the roots of a rose (Roza spec.) in a confined space**

In this research application the oxygen consumption of plant roots is measured by placing the roots of a plant in a confined container with a known volume sealed hermetically around the stem in order to avoid gas exchange of the confined container and surrounding. The oxygen consumption profile of the plants under different growth conditions can be easily determined.

Figure 8 shows a schematic representation of the experimental set-up of a plant (for example a rose) in a confined container.

Figure 9 shows the result. In this example two different kinds of metabolisms were found, depending on the oxygen concentration.

3. **Comparison of the start of germination between lettuce and barley seed**

In this set-up the germination speed of Lettuce (Grand Rapids Ritsa) seeds and Barley (Triumph 1989) seeds was compared in several containers. The seeds were confined in small confined containers with a volume of 200 microliter. The containers were scanned for their oxygen content every 30 minutes. By means of the measured oxygen concentration profile the start of germination for each species was calculated as shown in Figure 10. The experimental conditions were identical to the germination experiment described in experiment 1.

Figure 7 shows a schematic set-up of the confined containers for this experiment. The use of small confined containers as in this example shows that a method according to the invention is easily miniaturised.

Figure 10 shows the result of the above described experiment. This result is in accordance with the visual determination of germination of the 2 tested seeds.

5 4. **Detection of oxygen consumption of a worm**

The oxygen consumption of different worms was also determined with the Non Invasive Oxygen Detection (NIOD) Method. The individual worms were put into a confined space of only 200 microliters, provided with a small
10 spot of the oxygen sensitive coating.

Figure 11 shows a confined space comprising a worm and Figure 12 shows the results of the oxygen consumption of 2 different kinds of worms. It was calculated that the wood worm of 93 mg consumed about 0.5 microgram
15 oxygen per minute. This information was used to optimise the non-toxic killing method of wood worms in houses.

Figure legends

Figure 1 Measurement principle of optical oxygen sensor.

- 5 Figure 2 A rough schematic representation of a set-up for measuring metabolic gas changes and/or rates.

Figure 3 A detailed schematic representation of oxygen sensor.

- 10 Figure 4 Oxygen consumption during the germination of Triumph 1989 seeds at 25 °C.

Figure 5 Oxygen consumption during the germination of seeds.

- 15 Figure 6 Regression calculation.

Figure 7 Example of an automated version of a method according to the invention.

- 20 Figure 8 Experimental set-up for part of a plant in a confined container

Figure 9 Oxygen consumption of the roots of a Rose in a confined space

- 25 Figure 10 Determination of the start of germination by oxygen consumption in a confined space.

Figure 11 Picture showing a confined space comprising a worm

Figure 12 The oxygen consumption of a worm

References

1. Bambot S.B. et al., Phase Fluorimetric Sterilizable Optical Oxygen Sensor, Biotechnology and Bioengineering, 43:1139-1145, 1994.
- 5 2. Cox, M.E., Bunn, B., Detection of Oxygen by fluorescence quenching, Applied Optics, 24, 2114—2120, 1985
3. Holst, G.A., Flox an oxygen-flux-measuring system using a phase modulation method to evaluate the oxygen dependent fluorescent lifetime, Sensor and Actuators B 29 (1995) 231-239
- 10 4. Meier, B et al., Novel oxygen sensor material based on a ruthenium bipyridyl complex encapsulated in zeolyte Y: dramatic differences in the efficiency of luminescence quenching by oxygen on going from surface-adsorbed to zeolite-encapsulated fluorophores, Sensor and Actuators B 29 (1995) 240-245.
- 15 5. Marazuele, M.D. et al., Luminescence lifetime Quenching of a Ruthenium (II) Polypyridyl Dye for Optical Sensing of Carbon Dioxide. Appl. Spectroscopy (1998), 52:1314-1320.
6. A. Draaijer, J.W. J.W. König, J.J.F. van Veen, Substraat voor het inbedden van zuurstof gevoelige kleurstof, Dutch patent application, application
20 number 1014464

Claims

1. A method for determining metabolic state or rate or a change therein of at least one organism or part thereof comprising placing said organism or part thereof in a confined container and measuring the concentration of a metabolic gas in said confined container to determine consumption or production of said gas by said organism or part thereof wherein said gas concentration is determined without essentially affecting the concentration of said gas in said confined container.
2. A method according to claim 1 wherein said confined container is essentially not opened during measurements.
3. A method according to claim 1 or 2 wherein said gas comprises oxygen.
4. A method according to anyone of claims 1 to 3 wherein said gas concentration is determined optically.
5. A method according to claim 4 wherein said gas concentration is determined by determining the fluorescence quenching of a fluorescent dye present in said confined container.
6. A method according to claim 5 wherein said dye is present in a gas permeable compound.
7. A method according to claim 6 wherein said compound comprises a hydrophobic polymer.
8. A method according to anyone of claims 5 to 7 wherein said dye is present in at least a part of an inner coating of said confined container.
9. A method according to anyone of claims 1 to 8 wherein said organism comprises a seed.
10. A method according to claim 9 wherein said change in metabolic rate denotes germination.
11. A method according to anyone of claims 1 to 8 wherein said organism comprises a micro-organism.

12. A method according to claim 11 wherein said change in metabolic rate denotes cell activity.
13. A method according to claim 11 wherein said change in metabolic rate denotes cell death.
- 5 14. A method according to claim 11 wherein said change denotes sporulation.
15. A method according to claim 11 wherein said change denotes microbial fermentation.
16. A method to determine or monitor a rate of seed development
10 comprising use of a method according to claim 9 or 10.
17. A seed batch monitored with a method according to claim 16.
18. A method to determine or monitor a rate of culture development of a cell- or tissue-culture comprising use of a method according to anyone of claims 11 to 15.
- 15 19. A cell- or tissue culture monitored with a method according to claim 18.
20. A method to determine or monitor processing of waste water comprising use of a method according to anyone of claims 1 to 8.




Fig. 1

2/12




Fig. 2

3/12




Fig. 3

4/12

Oxygen consumption during the germination
of Triumph 1989 seeds at 25 °C




Fig. 4

5/12

Oxygen demand during the development of Triumph 1989 seeds at 25 °C





Fig. 5



Hours	2 (1 seed)	4 (1 seed)	5 (2 seeds)	6 (3 seeds)
6.5	277	275	253	227
7.5	271	279	231	204
8.5	282	287	257	198
9.5	269	279	227	160
11.5	275	284	202	125
14.0	262	268	172	
24.0	220	244		
29.0	250	263		
32.0	238	275		
35.5	226	267		
38.0	201	231		

Hours	2 (1 seed)	4 (1 seed)	5 (2 seeds)	6 (3 seeds)
6.5	0	0	23	56
7.5	2	4	26	52
8.5	8	0	48	75
9.5	-3	-8	22	81
11.5	10	0	52	119
14.0	4	-5	77	154
17	11	11	107	
24.0	59	35		
29.0	29	16		
32.0	41	4		
35.5	53	12		
38.0	78	48		

r.c.	2.001	0.950	9.948	17.747
Intercept	-12.083	-7.808	-38.721	-55.444
start hour	6.0	8.2	3.9	3.1

Fig. 6

7/12




Fig. 7




Fig. 8

9/12




Fig. 9

10/12

Determination of the start of the germination
by oxygen consumption in a confined space




Fig. 10

11/12




Fig. 11

12/12




Fig. 12

INTERNATIONAL SEARCH REPORT

Inte. .onal Application No

PCT/NL 01/00217

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 G01N33/497 A01C1/02 C12M1/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 G01N A01C C12M

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 54354 A (UNIV PENNSYLVANIA) 3 December 1998 (1998-12-03)	1-5, 11, 12, 15, 18-20
Y	the whole document	6, 7, 9, 10, 16, 17, 20
X	WO 98 12348 A (BECTON DICKINSON CO ;BURRELL GREGORY J (US); HU JOANNA KWOK YU (US) 26 March 1998 (1998-03-26) the whole document	1-5, 11, 12, 18, 19
Y	US 5 863 460 A (SLOVACEK RUDOLF E ET AL) 26 January 1999 (1999-01-26) abstract; figures	6, 7
	--- -/-- ---	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

10 May 2001

Date of mailing of the international search report

21/05/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bosma, R

INTERNATIONAL SEARCH REPORT

Inte. Jional Application No

PCT/NL 01/00217

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 36875 A (TEMPLETON COLIN W G ;COLOMBO STEPHEN J (CA)) 21 November 1996 (1996-11-21) page 22, line 6 -page 26, line 4 ---	9,10,16, 17
Y	AN L ET AL: "A NEW BIOSENSOR FOR RAPID OXYGEN DEMAND MEASUREMENT" WATER ENVIRONMENT RESEARCH,US,WATER ENVIRONMENT FEDERATION, ALEXANDRIA, vol. 70, no. 5, 1 July 1998 (1998-07-01), pages 1070-1074, XP000793304 ISSN: 1061-4303 page 1073, paragraph 3 ---	20
A	COX M E ET AL: "DETECTION OF OXYGEN BY FLUORESCENCE QUENCHING" APPLIED OPTICS,US,OPTICAL SOCIETY OF AMERICA,WASHINGTON, vol. 24, no. 14, 1985, pages 2114-2120, XP000885259 ISSN: 0003-6935 cited in the application the whole document ---	1-20
A	GIDROL X ET AL: "ACCUMULATION OF REACTIVE OXYGEN SPECIES AND OXIDATION OF CYTOKININ IN GERMINATING SOYBEAN SEEDS" EUROPEAN JOURNAL OF BIOCHEMISTRY,DE,BERLIN, vol. 224, no. 1, 15 August 1994 (1994-08-15), pages 21-28, XP000610708 ISSN: 0014-2956 abstract ---	9,10
A	EP 0 448 923 A (AVL PHOTRONICS CORP) 2 October 1991 (1991-10-02) abstract; figure 1 ---	1
A	DATABASE WPI Week 199213 Derwent Publications Ltd., London, GB; AN 1992-103639 XP002147978 & SU 1 645 893 A (REVUT V I), 30 April 1991 (1991-04-30) abstract -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 01/00217

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9854354 A	03-12-1998	US 6165741 A AU 7803098 A EP 0988396 A	26-12-2000 30-12-1998 29-03-2000
WO 9812348 A	26-03-1998	AU 4483997 A EP 1021557 A	14-04-1998 26-07-2000
US 5863460 A	26-01-1999	AU 2039597 A EP 0891540 A WO 9737210 A US 6074607 A	22-10-1997 20-01-1999 09-10-1997 13-06-2000
WO 9636875 A	21-11-1996	AU 5641996 A CA 2149508 A CA 2195108 A	29-11-1996 16-11-1996 21-11-1996
EP 0448923 A	02-10-1991	AT 132537 T DE 69024631 D DE 69024631 T US 5372936 A	15-01-1996 15-02-1996 19-09-1996 13-12-1994
SU 1645893 A	30-04-1991	NONE	

